

# Glossary

## COMP6741: Parameterized and Exact Computation

2017, Semester 2

### Glossary

**acyclic:** A graph is acyclic if it has no **cycle** as a **subgraph**.

**bipartite:** A graph  $G = (V, E)$  is bipartite if its vertex set can be partitioned into two **independent sets**. A partition  $(A, B)$  of  $V$  into independent sets is called a bipartition of  $G$ . The graph  $G$  is then often denoted by  $G = (A \uplus B, E)$ .

**Boolean formula:** A Boolean formula is constructed from Boolean variables that can take the values true and false (or 1 and 0) by the following operations: conjunction (AND,  $\wedge$ ), disjunction (OR,  $\vee$ ), and negation (NOT,  $\neg$ ).

**clique:** A subset of vertices  $S \subseteq V$  of a graph  $G = (V, E)$  is a clique in  $G$  if  $G[S]$  is a **complete** graph.

**closed neighborhood:** The closed neighborhood of a vertex  $v$  in a graph  $G$  is  $N_G[v] := \{v\} \cup N_G(v)$ . The subscript may be omitted if  $G$  is clear from the context.

**closed set neighborhood:** The closed neighborhood of a subset of vertices  $S \subseteq V$  in a graph  $G = (V, E)$  is  $N_G[S] := \bigcup_{v \in S} N_G[v]$ . The subscript may be omitted if  $G$  is clear from the context.

**coloring:** A coloring of a graph  $G = (V, E)$  is a function from  $V$  to a set of colors (integers) such that every two adjacent vertices in  $G$  are mapped to different colors. A  $k$ -coloring is a coloring using exactly  $k$  colors.

**complete:** A graph  $G$  is complete if there is an edge between each pair of vertices in  $G$ . A complete graph on  $n$  vertices is denoted by  $K_n$ .

**Conjunctive Normal Form:** A **Boolean formula** is in Conjunctive Normal Form if it is a conjunction of clauses, each clause is a disjunctions of literals, and each literal is a Boolean variable or its negation..

**connected:** A graph  $G$  is connected if there is a **walk** between every two vertices of  $G$ .

**connected component:** **Maximal connected subgraph**.

**cycle:** **2-regular connected** graph. A cycle on  $n$  vertices is denoted  $C_n$ .

**degree:** The degree of a vertex  $v$  in a graph  $G$  is  $d_G(v) := |N_G(v)|$ . The subscript may be omitted if  $G$  is clear from the context. The degree of a vertex  $v$  in a **multigraph:**  $G$  is the number of times  $v$  appears as an end point of an edge in  $E$ .

**directed acyclic graph:** A directed acyclic graph (DAG) is a **directed graph** that contains no **directed cycle** as a directed subgraph.

**directed cycle:** **Orientation** of a **cycle** where each vertex has in-degree 1.

**directed graph:** A directed graph  $G$  is an ordered pair  $(V, A)$  of a set  $V$  of vertices and a set  $A$  of arcs, where  $A$  is a set of ordered pairs of vertices. Its vertex set is  $V(G) = V$  and its arc set is  $A(G) = A$ .

**directed path:** **Orientation** of a **path** where each vertex has **in-degree** 1, except the start vertex, which has **in-degree** 0 and **out-degree** 1.

**disjoint union:** For two graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  with  $V_1 \cap V_2 = \emptyset$ , the disjoint union of  $G_1$  and  $G_2$ , denoted  $G_1 \oplus G_2$  is the graph  $(V_1 \cup V_2, E_1 \cup E_2)$ ; in case  $V_1 \cap V_2 \neq \emptyset$ , vertices need to be renamed before we can take the disjoint union of these graphs.

**distance:** In a graph  $G$ , the distance between two vertices  $u \in V$  and  $v \in V$  is the length of the shortest **walk** minus one between  $u$  and  $v$ , that is the minimum number of edges needed to be traversed to reach  $v$  from  $u$  and it is denoted by  $dist_G(u, v)$ .

**dominating set:** A subset of vertices  $S \subseteq V$  of a graph  $G = (V, E)$  is a dominating set of  $G$  if  $N_G[S] = V$ .

**feedback vertex set:** A feedback vertex set of a graph  $G = (V, E)$  is a subset of vertices  $S \subseteq V$  such that  $G - S$  is acyclic.

**forest:** **Acyclic** graph.

**graph:** A (simple, undirected) graph  $G$  is an ordered pair  $(V, E)$  of a set  $V$  of vertices and a set  $E$  of edges, where  $E$  is a set of unordered pairs of distinct vertices. Its vertex set is  $V(G) = V$  and its edge set is  $E(G) = E$ .

**in-degree:** The in-degree of a vertex  $v$  in a directed graph  $D$  is  $d_D^-(v) := |\{uv \in A\}|$ . The subscript may be omitted if  $D$  is clear from the context.

**independent set:** A subset of vertices  $S \subseteq V$  of a graph  $G = (V, E)$  is an independent set of  $G$  if  $G[S]$  has no edges.

**induced subgraph:** For a graph  $G = (V, E)$  and a vertex set  $S \subseteq V$ , the subgraph of  $G$  induced on  $S$  is the graph  $G[S] := (S, \{uv \in E : u, v \in S\})$ .

**maximal (set):** For a set  $\mathcal{S}$  of subsets of a ground set  $U$ , a set  $X \in \mathcal{S}$  is maximal if there exists no set  $Y \in \mathcal{S}$  with  $X \subsetneq Y$ .

**maximum (set):** For a set  $\mathcal{S}$  of subsets of a ground set  $U$ , a set  $X \in \mathcal{S}$  is maximum if there exists no set  $Y \in \mathcal{S}$  with  $|Y| > |X|$ .

**maximum degree:** The maximum degree of a graph  $G = (V, E)$  is  $\Delta(G) := \max_{v \in V} d_G(v)$ .

**minimum degree:** The minimum degree of a graph  $G = (V, E)$  is  $\delta(G) := \min_{v \in V} d_G(v)$ .

**multigraph:** A multigraph  $G$  is an ordered pair  $(V, E)$  of a set  $V$  of vertices and a *multiset*  $E$  of edges, where  $E$  is a multiset of unordered pairs of vertices. Its vertex set is  $V(G) = V$  and its edge set is  $E(G) = E$ .

**open neighborhood:** The (open) neighborhood of a vertex  $v$  in a graph  $G = (V, E)$  is  $N_G(v) := \{u \in V : uv \in E\}$ . The subscript may be omitted if  $G$  is clear from the context.

**open set neighborhood:** The (open) neighborhood of a subset of vertices  $S \subseteq V$  in a graph  $G = (V, E)$  is  $N_G(S) := N_G[S] \setminus S$ . The subscript may be omitted if  $G$  is clear from the context.

**order of growth:** Let  $g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  be a function. The set  $O(g(n))$  contains every function  $f$  such that there exist  $c, n_0 \geq 0$  such that  $f(n) \leq c \cdot g(n)$  for every  $n \geq n_0$ . The set  $o(g(n))$  contains every function  $f$  such that for every  $\epsilon > 0$  there exists a  $n_0 \geq 0$  such that  $f(n) \leq \epsilon \cdot g(n)$  for every  $n \geq n_0$ . For the set  $\Omega(g(n))$ , we have that  $f(n) \in \Omega(g(n))$  iff  $g(n) \in O(f(n))$ . For the set  $\omega(g(n))$ , we have that  $f(n) \in \omega(g(n))$  iff  $g(n) \in o(f(n))$ . For the set  $\Theta(g(n))$ , we have that  $\Theta(g(n)) = O(g(n)) \cap \Omega(g(n))$ .

**orientation:** An orientation of a graph  $G$  is a directed graph  $D$  that has exactly one arc for each edge of  $G$  with the same endpoints.

**out-degree:** The out-degree of a vertex  $v$  in a directed graph  $D$  is  $d_D^+(v) := |\{vu \in A\}|$ . The subscript may be omitted if  $D$  is clear from the context.

**path:** Tree with maximum degree 2. A path on  $n$  vertices is denoted  $P_n$ .

**regular:** A graph is  $d$ -regular if each of its vertices has degree  $d$ . A graph is regular if it is  $d$ -regular for some  $d$ .

**subgraph:** A graph  $H$  is a subgraph of a graph  $G$  if  $V(H) \subseteq V(G)$  and  $E(H) \subseteq E(G)$ .

**tree:** Acyclic, connected graph.

**vertex cover:** A subset of vertices  $S \subseteq V$  of a graph  $G = (V, E)$  is a vertex cover of  $G$  if each edge of  $G$  is incident to at least one vertex of  $S$ .

**vertex removal:** For a graph  $G = (V, E)$  and a vertex set  $S \subseteq V$ , the graph obtained by removing  $S$  from  $G$  is  $G - S := G[V \setminus S]$ . If  $S = \{u\}$ , we may write  $G - u$  instead of  $G - \{u\}$ .

**walk:** Sequence of vertices in a graph, with each vertex being adjacent to the vertices immediately preceding and succeeding it in the sequence.

## Problem Definitions

### $k$ -COLORING

Given a graph  $G$ , determine if there is a coloring of  $G$  with at most  $k$  colors.

### $k$ -SAT

Given a Boolean formula in Conjunctive Normal Form where each clause has at most  $k$  literals, determine if there is an assignment of its variables such that the formula evaluates to true.

### DOMINATING SET

Given a graph  $G$  and an integer  $k$ , determine whether  $G$  has a dominating set of size  $k$ .

### FEEDBACK VERTEX SET

Given a (multi)graph  $G$  and an integer  $k$ , determine whether  $G$  has a feedback vertex set of size at most  $k$ .

### INDEPENDENT SET

Given a graph  $G$  and an integer  $k$ , determine whether  $G$  has an independent set of size  $k$ .

#### MAXIMUM INDEPENDENT SET

Given a graph  $G$ , find an **independent set** of  $G$  of maximum cardinality.

#### MINIMUM VERTEX COVER

Given a graph  $G$ , find a **vertex cover** of  $G$  of minimum cardinality.

#### SAT

Given a Boolean formula, determine if there is an assignment of its variables such that the formula evaluates to **true**.

#### TRAVELING SALESMAN PROBLEM

Given a set  $\{1, \dots, n\}$  of  $n$  cities, the distance  $d(i, j)$  between every two cities  $i$  and  $j$ , and an integer  $k$ , determine whether there is a tour with total distance at most  $k$ . A *tour* is a permutation of the cities starting and ending in city 1.

#### VERTEX COVER

Given a graph  $G$  and an integer  $k$ , determine whether  $G$  has a **vertex cover** of size  $k$ .